

\$%^STN;HighlightOn= ***;HighlightOff=*** ;

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1600RKK

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS	1		Web Page for STN Seminar Schedule - N. America
NEWS	2	JUL 28	CA/CAPplus patent coverage enhanced
NEWS	3	JUL 28	EPFULL enhanced with additional legal status information from the epline Register
NEWS	4	JUL 28	IFICDB, IFIPAT, and IFIUDB reloaded with enhancements
NEWS	5	JUL 28	STN Viewer performance improved
NEWS	6	AUG 01	INPADOCDB and INPAFAMDB coverage enhanced
NEWS	7	AUG 13	CA/CAPplus enhanced with printed Chemical Abstracts page images from 1967-1998
NEWS	8	AUG 15	CAOLD to be discontinued on December 31, 2008
NEWS	9	AUG 15	CAPplus currency for Korean patents enhanced
NEWS	10	AUG 27	CAS definition of basic patents expanded to ensure comprehensive access to substance and sequence information
NEWS	11	SEP 18	Support for STN Express, Versions 6.01 and earlier, to be discontinued
NEWS	12	SEP 25	CA/CAPplus current-awareness alert options enhanced to accommodate supplemental CAS indexing of exemplified prophetic substances
NEWS	13	SEP 26	WPIDS, WPINDEX, and WPIX coverage of Chinese and Korean patents enhanced
NEWS	14	SEP 29	IFICLS enhanced with new super search field
NEWS	15	SEP 29	EMBASE and EMBAL enhanced with new search and display fields
NEWS	16	SEP 30	CAS patent coverage enhanced to include exemplified prophetic substances identified in new Japanese-language patents
NEWS	17	OCT 07	EPFULL enhanced with full implementation of EPC2000
NEWS	18	OCT 07	Multiple databases enhanced for more flexible patent number searching
NEWS	19	OCT 22	Current-awareness alert (SDI) setup and editing enhanced
NEWS	20	OCT 22	WPIDS, WPINDEX, and WPIX enhanced with Canadian PCT Applications
NEWS	21	OCT 24	CHEMLIST enhanced with intermediate list of pre-registered REACH substances
NEWS	22	NOV 21	CAS patent coverage to include exemplified prophetic substances identified in English-, French-, German-, and Japanese-language basic patents from 2004-present
NEWS	23	NOV 26	MARPAT enhanced with FSORT command
NEWS	24	NOV 26	MEDLINE year-end processing temporarily halts availability of new fully-indexed citations
NEWS	25	NOV 26	CHEMSAFE now available on STN Easy

NEWS 26 NOV 26 Two new SET commands increase convenience of STN
searching
NEWS 27 DEC 01 ChemPort single article sales feature unavailable
NEWS EXPRESS JUNE 27 08 CURRENT WINDOWS VERSION IS V8.3,
AND CURRENT DISCOVER FILE IS DATED 23 JUNE 2008.

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS LOGIN Welcome Banner and News Items
NEWS IPC8 For general information regarding STN implementation of
IPC 8

Enter NEWS followed by the item number or name to see news on that
specific topic.

All use of STN is subject to the provisions of the STN Customer
agreement. Please note that this agreement limits use to scientific
research. Use for software development or design or implementation
of commercial gateways or other similar uses is prohibited and may
result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 10:46:47 ON 02 DEC 2008

=> file agricola biosis embase caplus
COST IN U.S. DOLLARS

	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.42	0.42

FILE 'AGRICOLA' ENTERED AT 10:47:37 ON 02 DEC 2008

FILE 'BIOSIS' ENTERED AT 10:47:37 ON 02 DEC 2008
Copyright (c) 2008 The Thomson Corporation

FILE 'EMBASE' ENTERED AT 10:47:37 ON 02 DEC 2008
Copyright (c) 2008 Elsevier B.V. All rights reserved.

FILE 'CAPLUS' ENTERED AT 10:47:37 ON 02 DEC 2008
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2008 AMERICAN CHEMICAL SOCIETY (ACS)

=> s pineapple and organogenesis
L1 27 PINEAPPLE AND ORGANOGENESIS

=> l1 duplicate remove
L1 IS NOT A RECOGNIZED COMMAND
The previous command name entered was not recognized by the system.
For a list of commands available to you in the current file, enter
"HELP COMMANDS" at an arrow prompt (=>).

=> duplicate remove l1
DUPLICATE PREFERENCE IS 'AGRICOLA, BIOSIS, CAPLUS'
KEEP DUPLICATES FROM MORE THAN ONE FILE? Y/(N):n
PROCESSING COMPLETED FOR L1
L2 20 DUPLICATE REMOVE L1 (7 DUPLICATES REMOVED)

=> d l2 1-20 ibib ab

L2 ANSWER 1 OF 20 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States

of America. It contains copyrighted materials. All rights reserved.

(2008) on STN
ACCESSION NUMBER: 2008:52660 AGRICOLA <<LOGINID::20081202>>
DOCUMENT NUMBER: IND44033962
TITLE: The same treatment for transgenic shoot
regeneration elicits the opposite effect in mature explants
from two closely related sweet orange (Citrus
sinensis (L.)

Osborne) genotypes.
AUTHOR(S): Rodr Uguéz, Ana; Cervera, Magdalena; Peris, Josep
Enric; Pe la, Leandro
AVAILABILITY: DNAL (QK725.P53)
SOURCE: Plant cell, tissue, and organ culture, 2008
Apr. Vol.

93, no. 1 p. 97-106
Publisher: Dordrecht : Springer Netherlands
ISSN: 0167-6857

NOTE: Includes references
DOCUMENT TYPE: Article; (ELECTRONIC RESOURCE)
FILE SEGMENT: Non-US
LANGUAGE: English

AB In citrus, production of mature transgenic plants belonging to different genotypes is an important biotechnological objective. In the present study, we tried to genetically transform and regenerate mature plants from the economically important Navelina sweet orange cultivar by using the procedure previously established for the genetically close ***Pineapple*** sweet orange variety. The use of BAP at 3 mg l (British pound) promoted efficient shoot ***organogenesis*** in ***Pineapple*** as expected, but not in Navelina. Furthermore, different effects were observed when the auxin 1-naphtalene acetic acid (NAA) was added to BAP-containing regeneration media. Although NAA addition at 0.5 mg l (British pound) enhanced cambial callus formation, number of shoots and their elongation in Navelina, the contrary effect was observed in ***Pineapple***. Moreover, transformation efficiency in Navelina rose from 0 to 3% but declined from 6 to 0% in ***Pineapple***, indicating that BAP and BAP + NAA exerted the opposite effect in transgenic shoot regeneration from two closely related cultivars. This suggests that small changes in the procedure could induce drastic alterations in regeneration and even increase the likelihood of obtaining transformants from

non-responsive genotypes. Moreover, the vigour of the starting plant material and the addition of kanamycin as selective agent were determining for the generation of mature sweet orange transgenic plants.

L2 ANSWER 2 OF 20 BIOSIS COPYRIGHT (c) 2008 The Thomson Corporation on STN

DUPLICATE 2

ACCESSION NUMBER: 2008:530273 BIOSIS <<LOGINID::20081202>>

DOCUMENT NUMBER: PREV200800530272

TITLE: Establishment of callus induction and shoot regeneration

system from axillary bud on Taiwanese edible ***pineapples***.

AUTHOR(S): Zhang, Ya-Yen; Hsu, Huei-Juan; Huang, Wen-Lii [Reprint

Author]

CORPORATE SOURCE: Natl Chiayi Univ, Dept Agron, Chiayi, Taiwan wlhuang@mail.ncyu.edu.tw

SOURCE: Taiwanese Journal of Agricultural Chemistry and Food Science, (APR 2008) Vol. 46, No. 2, pp. 49-56. ISSN: 1605-2471.

DOCUMENT TYPE: Article

LANGUAGE: Chinese

ENTRY DATE: Entered STN: 24 Sep 2008

Last Updated on STN: 24 Sep 2008

AB Two cultivars, Tainung 17 (TNG-17) and Tainung 20 (TNG-20), of Taiwanese

edible ***pineapple*** (Ananas comosus L. Merrill) were used in this

study. The axillary buds from ***pineapple*** crown grown in the

field were selected and inoculated on MS basal medium supplement with

different combinations of NAA and BA. It showed that the callus could be

induced when NAA supplemented in the medium. However, the texture is

loose and browning. Besides, protocorm-like body (PLB) formed when the

explants were inoculated on the medium containing BA and NAA.

After being

transferred the callus and PLB onto MSB4N4 medium, somatic embryogenesis

will be mainly observed in TNG-20. However, somatic embryogenesis and

organogenesis are observed in TNG-17 during cell differentiation.

We have developed efficient methods for plant regeneration, via both

embryogenesis and ***organogenesis*** on Taiwanese edible ***pineapple***.

In addition, we also found abundance of

starch granules spread in TNG-17 callus. It is the first discovery in

pineapple tissue culture. Further studies is necessary to

illuminate the possible roles of starch accumulation during callus induction and cell differentiation.

L2 ANSWER 3 OF 20 BIOSIS COPYRIGHT (c) 2008 The Thomson Corporation
on STN

ACCESSION NUMBER: 2008:374461 BIOSIS <<LOGINID::20081202>>
DOCUMENT NUMBER: PREV200800374460
TITLE: Protocols for Micropropagation of Woody Trees and
Fruits.
AUTHOR(S): Jain, SM [Editor]; Haggman, H [Editor]
SOURCE: Jain, SM [Editor]; Haggman, H [Editor]. (2007)
Protocols

for Micropropagation of Woody Trees and Fruits.
Publisher: SPRINGER, PO BOX 17, 3300 AA DORDRECHT,
NETHERLANDS.
ISBN: 978-1-4020-6351-0(H).

DOCUMENT TYPE: Book
LANGUAGE: English
ENTRY DATE: Entered STN: 2 Jul 2008
Last Updated on STN: 2 Jul 2008

AB This 544-page book is a summary of protocols for micropropagation
of woody
trees and fruits. There are 48-individually authored chapters
organized
in three sections. The first section deals with totipotency, cell
cycle,
micropropagation via ***organogenesis*** in slash pine,
micropropagation of Sequoia sempervirens, Pinus pinea, Pinus
armandii var.
Amamiana, ***organogenesis*** and cryopreservation of juvenile
radiata
pine, genetic fidelity analyses, micropropagation of Quercus,
Cupressus
sempervirens, Taxus baccata and propagation of selected Pinus
genotypes,
protocol for doubled-haploid micropropagation, in vitro propagation
of
Fraxinus species and Ulmus species. The other topics include
micrografting, in vitro conservation and micropropagation in
grapevine and
pistachio, in vitro mutagenesis and mutant multiplication,
micropropagation protocol for microspore embryogenesis in Olea
europaea,
tissue culture propagation and high frequency shoot formation
protocol,
micropropagation of selected Vaccinium species, ***pineapple***
,
micropropagation of bamboo species through axillary shoot
proliferation
and light-emitting diodes as an effective lighting source for in
vitro
banana culture. The text is written in English, followed by a set
of
references at the end of each chapter. Users of this book will
include
graduate students and researchers in plant tissue culture and
micropropagation.

L2 ANSWER 4 OF 20 CAPLUS COPYRIGHT 2008 ACS on STN
ACCESSION NUMBER: 2008:107748 CAPLUS <<LOGINID::20081202>>
DOCUMENT NUMBER: 148:208004
TITLE: Medicinal plant biotechnology research in
Jamaica -

challenges and opportunities
 AUTHOR(S): Mitchell, S. A.; Ahmad, M. H.
 CORPORATE SOURCE: Medicinal Plant Research Group, Biotechnology
 Centre,
 University of the West Indies, West Indies,
 Jamaica
 SOURCE: Acta Horticulturae (2007), 756 (Proceedings of
 the
 International Symposium on Medicinal and
 Nutraceutical
 Plants, 2007), 171-181
 CODEN: AHORA2; ISSN: 0567-7572
 PUBLISHER: International Society for Horticultural Science
 DOCUMENT TYPE: Journal; General Review
 LANGUAGE: English
 AB A review on the authors' own work. Medicinal Plant Biotechnol.
 Research
 in a tropical developing country is a challenge but there are many
 opportunities as well. This paper reviews research undertaken by
 the
 Medicinal Plant Research Group from its inception in 1999 to the
 end of
 2006. A three-prong approach has been taken to maintain an
 international
 std. of research while ensuring local and regional relevancy: 1)
 formulation of antimicrobial products (including a neem *Azadirachta*
indica) disinfectant); 2) tissue culture studies (micropropagation
 of
 medicinal plants including neem, ginger [*Zingiber officinalis*],
 turmeric
 [*Curcuma longa*], leaf-of-life [*Bryophyllum pinnatum*], Quako
 [*Mikania*
micrantha], John Charles [*Hyptis verticillata*], peperomia
 [*Peperomia*
hernandifolia], nail cleaner [*Arthrostema fragile*], lemon grass
 [*Cymbopogon citratus*], ***pineapple*** [*Ananas cosmosus*] and
 sarsaparilla [*Smilax regelii*]; somatic embryogenesis of ackee
 [*Blighia*
sapida] and guinea hen weed [*Petiveria alliacea*]; and de novo
 organogenesis of scotch bonnet pepper [*Solanum*
chinense]); and 3)
 business studies including information gathering and dissemination
 (Jamaican folk medicine practises, UWI medicinal plant research
 1948-2001,
 e-book on Caribbean medicinal plants, book chapter on Jamaica's
 medicinal
 plant biotechnol. experience, article on medicinal gene bank and
 folk
 recipes of 30 of these plants, over 56 newspaper articles, 13
 e-newsletters, marketing and feasibility studies and business
 plans, plus
 several presentations at various audiences including of scientists,
 farmers, government bodies and industrial groups). There has been
 a
 conscious effort to be involved in and to tailor research to serve
 industrial needs. There has also been a conscious effort to mix
 short-term research that has immediate application (eg. development
 of
 low-cost tissue culture kits) with longer-term research that may
 take

years to apply but for which the potential is much greater (e.g.,
mol.

pharming, and somatic embryogenesis of elite trees). The
challenges and
opportunities arising from these activities will be discussed.

REFERENCE COUNT: 39 THERE ARE 39 CITED REFERENCES AVAILABLE
FOR THIS

RECORD. ALL CITATIONS AVAILABLE IN THE RE
FORMAT

L2 ANSWER 5 OF 20 BIOSIS COPYRIGHT (c) 2008 The Thomson Corporation
on STN

ACCESSION NUMBER: 2007:308300 BIOSIS <<LOGINID::20081202>>

DOCUMENT NUMBER: PREV200700291509

TITLE: Callogenesis and ***organogenesis*** in
pineapple : a histological and

ultrastructural study

of developing callus and morphogenic processes.

AUTHOR(S): Bennici, A. [Reprint Author]; Mori, B.; Tani, C.;
Bussi, B.

CORPORATE SOURCE: Univ Florence, Dipartimento Biol Vegetale, Piazzale
Cascine

28, I-50144 Florence, Italy

SOURCE: Advances in Horticultural Science, (2007) Vol. 21,
No. 1,

pp. 19-27.

ISSN: 0394-6169.

DOCUMENT TYPE: Article

LANGUAGE: English

ENTRY DATE: Entered STN: 9 May 2007

Last Updated on STN: 9 May 2007

AB Organogenic callus cultures from young leaf explants of Ananas
comosus

(L.) Merr. var. Smooth Cayenne cv. Serrana were obtained on
Murashige

and Skoog (1962) medium using eight different protocols with regard
to the

growth regulator types and/or combinations and doses tested
(dicamba and

benzyladenine, picloram or 2,4-dichlorophenoxyacetic acid and
benzyladenine, dicamba and kinetin). In some cases, "shoot

inducing

medium" containing BA or kinetin alone, after a "callus inducing
medium",

were also used. The various media tested did not influence the
number of

explants forming callus (practically 100%) and the growth of the
calluses,

as well as the type of organogenic process (shoot regeneration) and
its

frequency, except for the medium containing 2,4-D. This compound
at 2.5

mg l(-1) doubled the-total I final callus mass, in comparison to
the other

media, and induced shoots or shoots and roots, whereas at 4.0 mg
l(-1)stimulated only root formation. Also the transfer of callus

to a

shoot-inducing medium did not significantly influence shoot
regeneration

rate. Light and electron microscope analysis showed similar

patterns of
callus formation in all the explants, where callus initiation
occurred
from parenchyma cells surrounding the vascular bundles. During
callus
growth meristematic centers appeared at its periphery. Thereafter,
they
developed into shoots (or roots). Roots were never associated
directly
with shoots. Plant regeneration was always by indirect
organogenesis and never the result of somatic
embryogenesis. The
meristematic and organogenic activity were found to be related to
abundant
starch and protein accumulation in the parenchyma cells located
under or
near the meristems, with vascular connections with the callus
itself, and
with a strong thickening of the walls around the meristematic
zones.

L2 ANSWER 6 OF 20 BIOSIS COPYRIGHT (c) 2008 The Thomson Corporation
on STN

DUPLICATE 3
ACCESSION NUMBER: 2006:569014 BIOSIS <<LOGINID::20081202>>
DOCUMENT NUMBER: PREV200600556509
TITLE: The introduction of transgenes to control blackheart
in
pineapple (Ananas comosus L.) cv. Smooth
cayenne by
microprojectile bombardment.
AUTHOR(S): Ko, H. L. [Reprint Author]; Campbell, P. R.; Jobin-
Decor,
M. P.; Eccleston, K. L.; Graham, M. W.; Smith, M. K.
CORPORATE SOURCE: Dept Primary Ind and Fisheries, Maroochy Res Stn,
SCMC, POB
5083, Nambour, Qld 4560, Australia
Lien.Ko@dpi.qld.gov.au
SOURCE: Euphytica, (AUG 2006) Vol. 150, No. 3, pp. 387-395.
CODEN: EUPHAA. ISSN: 0014-2336.
DOCUMENT TYPE: Article
LANGUAGE: English
ENTRY DATE: Entered STN: 27 Oct 2006
Last Updated on STN: 27 Oct 2006

AB A transformation technique for the introduction of transgenes to
control
blackheart by particle bombardment has been developed for
pineapple cv. Smooth Cayenne. Leaf callus cultures
capable of
high frequency ***organogenesis*** with a short regeneration
time were
used as explant material. Gus and gfp reporter genes were used to
observe
and determine transient and stable expression. The ppo gene,
isolated
from ***pineapple***, was introduced to control blackheart.
Co-transformation occurred with constructs containing the nptII
gene
conferring geneticin resistance. We have recovered 15 independent
transgenic gus and gfp lines each from 8 separate experiments and

22 ppo
lines from 11 experiments. Gus, gfp, ppo and nptII positive plants
have
been regenerated, which have been shown by Southern blot analysis
to be
stable transgenics containing multiple copies of the introduced
genes.
These results show that biolistic gene delivery in
pineapple can
be successfully achieved at an acceptable efficiency of 0.21-1.5%
for
genetic improvement of 'Smooth Cayenne', the industry standard
throughout
the world.

L2 ANSWER 7 OF 20 AGRICOLA Compiled and distributed by the National
Agricultural Library of the Department of Agriculture of the United
States
of America. It contains copyrighted materials. All rights
reserved.

(2008) on STN DUPLICATE 4
ACCESSION NUMBER: 2006:68161 AGRICOLA <<LOGINID::20081202>>
DOCUMENT NUMBER: IND43828493
TITLE: Transformation and regeneration of
pineapple

AUTHOR(S): Firoozabady, E.; Heckert, M.; Gutterson, N.
SOURCE: Plant cell, tissue and organ culture, 2006 Jan.
Vol.

84, no. 1 p. 1-16
ISSN: 0167-6857

NOTE: Includes references
DOCUMENT TYPE: Article; (ELECTRONIC RESOURCE)
FILE SEGMENT: Non-US
LANGUAGE: English

AB We have developed efficient methods for plant regeneration, via
both

organogenesis and embryogenesis, of Smooth Cayenne
pineapple , Ananas comosus (L.) Merr. A range of different
types of

embryogenic tissues has been developed with varying properties in
terms of
growth rate and state of development (Firoozabady and Moy, 2004).

Two of
the embryogenic systems, namely friable embryogenic cell clusters
(ECCs)
and chunky non-dispersible embryogenic tissues (ETs) have been used
for

transformation of ***pineapple*** . The tissues were
cocultivated for

2-3 days with Agrobacterium tumefaciens disarmed strain C58
carrying a
binary vector containing either surB gene conferring resistance to
chlorsulfuron or the nptII gene conferring resistance to geneticin
(G418).

After cocultivation and a recovery period, tissues were selected on
media

containing chlorsulfuron or G418. On average, about 50 or 120
independent

transgenic lines were obtained from each gram of ECCs or ETs,

respectively, inoculated with Agrobacterium. Transformed embryogenic tissues were transferred to maturation media to form somatic embryos, which subsequently produced transgenic ***pineapple*** plants. Transformation has been confirmed by GUS assay, polymerase chain reaction, and by Southern hybridization. Thousands of plants from independently transformed lines were transferred to the greenhouse and to the field to evaluate clonal fidelity and somaclonal variation.

L2 ANSWER 8 OF 20 BIOSIS COPYRIGHT (c) 2008 The Thomson Corporation on STN

DUPLICATE 5

ACCESSION NUMBER: 2006:321571 BIOSIS <<LOGINID::20081202>>

DOCUMENT NUMBER: PREV200600317177

TITLE: Glutamine enhances competence for

organogenesis

in ***pineapple*** leaves cultivated in vitro.

AUTHOR(S): Hamasaki, Regina M.; Purgatto, Eduardo; Mercier, Helenice

[Reprint Author]

CORPORATE SOURCE: Univ Sao Paulo, Dept Bot, CP 11461, BR-05422970 Sao Paulo,

SP, Brazil

hmercier@usp.br

SOURCE: Brazilian Journal of Plant Physiology, (OCT-DEC 2005) Vol.

17, No. 4, pp. 383-389.

ISSN: 1677-0420.

DOCUMENT TYPE: Article

LANGUAGE: English

ENTRY DATE: Entered STN: 21 Jun 2006

Last Updated on STN: 21 Jun 2006

AB Leaf bases of ***pineapple*** cultured on a shoot induction medium

(SIM) produced protuberances followed by shoot-buds via direct

organogenesis at a frequency of 46 %. When 8 mM

glutamine (gin)

was a supplement to SIM (SIM8gln), the regeneration rate increased to 70

%, thus suggesting that 8mM gin increased explant competence for

organogenesis. Besides this, shoot vigor was strongly

enhanced in

SIM8gln. Other gin concentrations (16 or 32 mM) evoked a lower frequency

of shoot-bud induction and number of regenerated shoots per explant when

compared to SIM8gln. In this study, it was defined that explant organogenic commitment to form shoot-buds occurred in the first 7

days of

culture on SIM8gln. Thereafter, endogenous indole-3-acetic acid (IAA) and

cytokinin (4 types) measurements were carried out during this period, that

is, during the induction phase of shoot-bud formation. The IAA content

increased greatly until the 5(th), day in the leaf bases cultured

on

SIM8gln. No such change in IAA concentration was observed in the explants

cultivated on SIM or in the presence of the highest gin concentration (32

mM), this being inhibitory to the organogenic process. The only natural

cytokinin detected was isopentenyladenine. An increase of 50 % in the

level of this phytohormone occurred in leaf bases cultured on SIM8gln at

the 5(th) day, when compared to SIM or of 170% compared to SIM32gln.

These results suggest that 8 mM gin favorably influenced the organogenic

process through changes in IAA and iP concentrations in ***pineapple***

leaves.

L2 ANSWER 9 OF 20 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States

of America. It contains copyrighted materials. All rights reserved.

(2008) on STN

DUPLICATE 6

ACCESSION NUMBER: 2004:47009 AGRICOLA <<LOGINID::20081202>>

DOCUMENT NUMBER: IND43646424

TITLE: Regeneration of ***pineapple*** plants in via

somatic embryogenesis and ***organogenesis***

AUTHOR(S): Firoozabady, E.; Moy, Y.

AVAILABILITY: DNAL (QK725.I43)

SOURCE: In vitro cellular & developmental biology - Plant,

2004 Jan-Feb Vol. 40, no. 1 p. 67-74

ISSN: 1054-5476

NOTE: Includes references

DOCUMENT TYPE: Article

FILE SEGMENT: Other US

LANGUAGE: English

L2 ANSWER 10 OF 20 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States

of America. It contains copyrighted materials. All rights reserved.

(2008) on STN

DUPLICATE 7

ACCESSION NUMBER: 2003:59403 AGRICOLA <<LOGINID::20081202>>

DOCUMENT NUMBER: IND23346548

TITLE: Plant regeneration by somatic embryogenesis and ***organogenesis*** in commercial

pineapple

(Ananas comosus L.).

AUTHOR(S): Sripaoraya, S.; Marchant, R.; Power, J.B.;

Davey, M.R.

AVAILABILITY: DNAL (QK725.I43)

SOURCE: In vitro cellular & developmental biology.

Plant :

journal of the Tissue Culture Association,

Sept/Oct

2003. Vol. 39, No. 5. p. 450-454
Publisher: Largo, MD : Society for In Vitro

Biology.

CODEN: IVCPEO; ISSN: 1054-5476

NOTE:

Includes references

PUB. COUNTRY:

Maryland; United States

DOCUMENT TYPE:

Article

LANGUAGE:

English

L2 ANSWER 11 OF 20 BIOSIS COPYRIGHT (c) 2008 The Thomson Corporation
on

STN

ACCESSION NUMBER: 2004:161532 BIOSIS <<LOGINID::20081202>>

DOCUMENT NUMBER: PREV200400159398

TITLE: Levels of endogenous free amino acids during
induction

phase of shoot ***organogenesis*** in leaves of
pineapple cultured in vitro.

AUTHOR(S):

Kitakawa, Adelia Y. [Reprint Author]; Hamasaki,

Regina M.

[Reprint Author]; Mercier, Helenice [Reprint Author]

CORPORATE SOURCE:
05422-970,

Department of Botany, Sao Paulo University, CEP

CP 11461, Sao Paulo, SP, Brazil

SOURCE:
2, pp.

Amino Acids (Vienna), (September 2003) Vol. 25, No.

170. print.

Meeting Info.: 8th International Congress on Amino

Acids

and Proteins. Rome, Italy. September 05-09, 2003.

ISSN: 0939-4451.

DOCUMENT TYPE:

Conference; (Meeting)

Conference; Abstract; (Meeting Abstract)

LANGUAGE:

English

ENTRY DATE:

Entered STN: 24 Mar 2004

Last Updated on STN: 24 Mar 2004

L2 ANSWER 12 OF 20 AGRICOLA Compiled and distributed by the National
Agricultural Library of the Department of Agriculture of the United
States

of America. It contains copyrighted materials. All rights
reserved.

(2008) on STN

ACCESSION NUMBER: 2003:59305 AGRICOLA <<LOGINID::20081202>>

DOCUMENT NUMBER: IND23346446

TITLE: Micropropagation of ***pineapple*** guava
through

organogenesis and axillary shoot
proliferation.

AUTHOR(S):

Canhoto, J.M.; Gruz, G.S.

AVAILABILITY:

DNAL (80 Ac82)

SOURCE:
117

Acta horticulturae, Jan 2000. No. 520. p. 109-

Publisher: Leuven, Belgium : International

Society for

Horticultural Science.

CODEN: AHORA2; ISSN: 0567-7572

NOTE:

Paper presented at the Twenty-fifth

International

Horticultural Congress, August 2-7, 1998,
 Brussels,
 Belgium. Part 10.
 Includes references
 PUB. COUNTRY: Belgium
 DOCUMENT TYPE: Article
 LANGUAGE: English

L2 ANSWER 13 OF 20 BIOSIS COPYRIGHT (c) 2008 The Thomson Corporation
 on
 STN
 ACCESSION NUMBER: 2000:242637 BIOSIS <<LOGINID::20081202>>
 DOCUMENT NUMBER: PREV200000242637
 TITLE: Auxin/cytokinin control of shoot
 organogenesis in
 pineapple leaf explants.
 AUTHOR(S): Mercier, H. [Reprint author]; Souza, B. M. [Reprint
 author]
 CORPORATE SOURCE: Department of Botany, University of Sao Paulo, Sao
 Paulo,
 Brazil
 SOURCE: Biologia Plantarum (Prague), (1999) Vol. 42, No.
 SUPPL.,
 pp. S53. print.
 Meeting Info.: International Symposium on Auxins and
 Cytokinins in Plant Development. Prague, Czech
 Republic.
 July 26-30, 1999. Institute of Experimental Botany,
 Academy
 of Sciences of the Czech Republic.
 CODEN: BPABAJ. ISSN: 0006-3134.
 DOCUMENT TYPE: Conference; (Meeting)
 Conference; Abstract; (Meeting Abstract)
 LANGUAGE: English
 ENTRY DATE: Entered STN: 14 Jun 2000
 Last Updated on STN: 5 Jan 2002

L2 ANSWER 14 OF 20 CAPLUS COPYRIGHT 2008 ACS on STN
 ACCESSION NUMBER: 1997:679171 CAPLUS <<LOGINID::20081202>>
 DOCUMENT NUMBER: 127:327456
 ORIGINAL REFERENCE NO.: 127:64169a,64172a
 TITLE: Regulated excision of a target gene from the
 transformation vector in the recipient cell
 using a
 site-specific recombinase
 INVENTOR(S): Surin, Brian Peter; De Feyter, Robert Charles;
 Graham,
 Michael Wayne; Waterhouse, Peter Michael;
 Keese, Paul
 Konrad; Shahjahan, Ali
 PATENT ASSIGNEE(S): Commonwealth Scientific and Industrial Research
 Organisation, Australia; The Australian
 National
 University; Surin, Brian Peter; De Feyter,
 Robert
 Charles; Graham, Michael Wayne; Waterhouse,
 Peter
 Michael; Keese, Paul Konrad; Shahjahan, Ali
 SOURCE: PCT Int. Appl., 85 pp.
 CODEN: PIXXD2

DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9737012	A1	19971009	WO 1997-AU197	
19970327				
W:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, ML, MR, NE, SN, TD, TG			
CA 2250111	A1	19971009	CA 1997-2250111	
19970327				
AU 9721437	A	19971022	AU 1997-21437	
19970327				
AU 717267	B2	20000323		
EP 922097	A1	19990616	EP 1997-913984	
19970327				
R:	BE, CH, DE, ES, FR, GB, IT, LI, NL, SE			
NZ 331940	A	20000228	NZ 1997-331940	
19970327				
JP 2000507446	T	20000620	JP 1997-534743	
19970327				
US 20020147168	A1	20021010	US 2001-850846	
20010507				
PRIORITY APPLN. INFO.:			AU 1996-9031	A
19960329			WO 1997-AU197	W

19970327
 AB A method of site-specific excision of a target gene from a transformation vector using a site-specific recombinase is described. This allows the transformation of the target organism with the removal of a selectable marker carried by the vector. Excision can be regulated or constitutive depending upon the promoter regulating the recombinase gene. As a result the same selectable marker can be used can be used in a no. of sequential transformations. The method can be generally used to regulate transgene expression in genetically-manipulated organisms, for example to promote differentiation, de-differentiation, or any unidirectional developmental

shift of a target cell which requires the time-specific expression of a particular gene. The method is particularly suited to the promotion of specific organogeneses in plants using ***organogenesis*** - promoting transgenes, wherein the organs which subsequently develop in said plants are genetically transformed with a desired gene but lack ***organogenesis*** -promoting transgenes. The use flp/frt and cre/loxP recombination systems in tobacco (*Nicotiana plumbaginifolia*) is demonstrated.

L2 ANSWER 15 OF 20 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1996:492484 CAPLUS <<LOGINID::20081202>>

DOCUMENT NUMBER: 125:137763

ORIGINAL REFERENCE NO.: 125:25681a,25684a

TITLE: Feijoa sellowiana Berg (***pineapple*** guava)

AUTHOR(S): Canhoto, J. M.; Cruz, G. S.

CORPORATE SOURCE: Departamento de Botanica, Universidade de Coimbra,

Coimbra, 3049, Port.

SOURCE: Biotechnology in Agriculture and Forestry (1996),

35(Trees IV), 155-171

CODEN: BAFOEG; ISSN: 0934-943X

PUBLISHER: Springer

DOCUMENT TYPE: Journal; General Review

LANGUAGE: English

AB A review with 30 refs. on somatic embryogenesis, shoot multiplication and

organogenesis studies of *F. sellowiana*.

L2 ANSWER 16 OF 20 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1995:612878 CAPLUS <<LOGINID::20081202>>

DOCUMENT NUMBER: 123:107960

ORIGINAL REFERENCE NO.: 123:19147a,19150a

TITLE: Effect of various media composition on in vitro propagation of *Ananas comosus* (L.) Merr.

AUTHOR(S): Bordoloi, Nabanita Dutta; Sarma, C. M.

CORPORATE SOURCE: Department Botany, Gauhati University, Gauhati, 781

014, India

SOURCE: Journal of Plant Science Research (1994), Volume Date

1993, 9(1-4), 50-3

CODEN: JPSREB; ISSN: 0970-2539

PUBLISHER: Society for the Promotion of Plant Science Research

DOCUMENT TYPE: Journal

LANGUAGE: English

AB In vitro micropropagation of ***pineapple*** , *Ananas comosus* (L.) Merr.

(cv. Queen) was studied in relation to various concns. of several PGRs,

sucrose, macro- and micronutrients. Four nutrient media viz. MS, B5, SH

and White's contg. various nutrients were tested for callus

formation,
 organogenesis , plantlet formation and development of
 roots. MS
 macronutrients with B5 micronutrients showed satisfactory results
 in both
 callus formation and ***organogenesis*** . Both MS and SH media
 supplemented with IAA, IBA and kinetin (KN) (5 .mu.g mL-1 each)
 exhibited
 good response in the establishment of solitary shoots. Profuse
 shoot
 formation was obsd. in MS medium supplemented with various concns.
 of IBA,
 KN and CH. Callus initiation at the base of the in vitro obtained
 shoot
 explants was also obsd. in MS medium supplemented with IAA, IBA and
 KN (5
 .mu.g mL-1 each). Regenerated shoots produced roots on both half
 strength
 salt of MS and B5 basal media supplemented with IBA or NAA (2 .mu.g
 mL-1).
 Rooted plantlets were transplanted to earthen pots filled with
 sterilized
 sand. The pots were regularly watered and nutrient solns. added.
 After
 acclimatization in the earthen pots, plantlets were transferred to
 the
 natural condition in the field. The rate of survival was 95-97%.

L2 ANSWER 17 OF 20 AGRICOLA Compiled and distributed by the National
 Agricultural Library of the Department of Agriculture of the United
 States

of America. It contains copyrighted materials. All rights
 reserved.

(2008) on STN

ACCESSION NUMBER:	91:81856 AGRICOLA <<LOGINID::20081202>>
DOCUMENT NUMBER:	IND91045962
TITLE:	Growth and morphogenesis of citrus tissue
cultures	infected with psorosis, vein enation, and
cachexia.	
AUTHOR(S):	Duran-Vila, N.; Medina, V.; Pina, J.A.; Ortega, C.;
	Molins, M.I.; Navarro, L.
CORPORATE SOURCE:	Instituto Valenciano de Investigaciones
Agrarias,	
	Valencia, Spain
AVAILABILITY:	DNAL (464.8 P56)
SOURCE:	Phytopathology, Aug 1991. Vol. 81, No. 8. p.
824-831	
	Publisher: St. Paul, Minn. : American
	Phytopathological Society.
	CODEN: PHYTAJ; ISSN: 0031-949X
NOTE:	Includes references.
DOCUMENT TYPE:	Article
FILE SEGMENT:	U.S. Imprints not USDA, Experiment or Extension
LANGUAGE:	English
AB Stem segments from	***Pineapple*** sweet orange (Citrus
sinensis) and	
	Etrog citron (C. medica) infected with psorosis, vein enation, and
	cachexia, as well as uninfected controls, were cultured in vitro.

Production of roots and regeneration of shoots and buds were modified as a result of infection. The number of explants showing morphogenesis and the amount of rooting and/or regeneration of shoots and buds were affected as compared with the uninfected explants cultured as controls. The differences on morphogenic patterns depended on the disease and the disease isolate. Explants infected with vein enation and cachexia produced significantly less primary callus than the controls, whereas psorosis did not affect callus induction. The amount and morphology of secondary callus after the first subculture were similar in infected and uninfected tissues. Biological indexing of callus indicated that psorosis- and cachexia-infected callus were good host systems for the replication of the disease-causing agents, whereas vein enation could not be detected after continuous callus cultures. The citrus cachexia viroid was detected from infected callus by nucleic acid extraction and sequential polyacrylamide gel electrophoresis. Electron microscopy studies revealed alterations at the cell level on psorosis-infected callus.

L2 ANSWER 18 OF 20 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States

of America. It contains copyrighted materials. All rights reserved.

(2008) on STN

ACCESSION NUMBER:	92:21141 AGRICOLA <<LOGINID::20081202>>
DOCUMENT NUMBER:	IND92003950
TITLE:	***Organogenesis*** in callus cultures of ***pineapple*** (Ananas comosus (L.)
Merr.).	
AUTHOR(S):	Fitchet, M.
CORPORATE SOURCE:	Citrus and Subtropical Fruit Research
Institute,	
	Nelspruit, Republic of South Africa
AVAILABILITY:	DNAL (80 AC82)
SOURCE:	Acta horticultrae, July 1990. No. 275. p. 267-
274	
	Publisher: Wageningen : International Society
for	
	Horticultural Science.
	CODEN: AHORA2; ISSN: 0567-7572
NOTE:	Paper presented at the "International Symposium
on the	
	Culture of Subtropical and Tropical Fruits and
Crops,"	
	Volume I, November 6-10, 1989, Nelspruit, South
	Africa.
	Includes references.
DOCUMENT TYPE:	Article
FILE SEGMENT:	Non-U.S. Imprint other than FAO
LANGUAGE:	English

AB Callus was induced from the crown apical region of 'Queen'
 pineapples on Murashige and Tucker medium with casein
 hydrolysate
 (400 mg/l), coconut water (15%) and naphthaleneacetic acid (40
 mg/l).
 Callus did not become organogenic unless it passed through a stage
 where
 the colour changed from yellow to green. By investigating the
 anatomical
 changes in the green callus it was possible to determine that the
 regeneration of plants was by indirect adventitious
 organogenesis
 , and not the result of somatic embryogenesis. Areas of
 meristematic
 activity were easily discernible, and developing shoot buds could
 be seen
 on the periphery of the callus as well as within the callus mass.

L2 ANSWER 19 OF 20 BIOSIS COPYRIGHT (c) 2008 The Thomson Corporation
 on

STN
 ACCESSION NUMBER: 1986:428160 BIOSIS <<LOGINID::20081202>>
 DOCUMENT NUMBER: PREV198631093972; BR31:93972
 TITLE: TISSUE CULTURE RESEARCH AT NATIONAL INSTITUTE OF
 AGROBIOLOGICAL RESOURCES JAPAN.
 AUTHOR(S): SHIGA T [Reprint author]
 CORPORATE SOURCE: DEP OF CELL BIOL, NATL INST OF AGROBIOLOGICAL
 RESOURCES,
 YATABE, TSUKUBA, IBARAKI, 305, JAPAN
 SOURCE: (1985) pp. 349-358. INTERNATIONAL RICE RESEARCH
 INSTITUTE.
 BIOTECHNOLOGY IN INTERNATIONAL AGRICULTURAL
 RESEARCH;
 INTER-CENTER SEMINAR ON INTERNATIONAL AGRICULTURAL
 RESEARCH
 CENTERS AND BIOTECHNOLOGY, MANILA, PHILIPPINES, APR.
 23-27,
 1984. VIII+435P. INTERNATIONAL RICE RESEARCH
 INSTITUTE:
 MANILA, PHILIPPINES. ILLUS. PAPER.
 ISBN: 971-104-124-3.
 DOCUMENT TYPE: Book
 Conference; (Meeting)
 FILE SEGMENT: BR
 LANGUAGE: ENGLISH
 ENTRY DATE: Entered STN: 25 Oct 1986
 Last Updated on STN: 25 Oct 1986

L2 ANSWER 20 OF 20 BIOSIS COPYRIGHT (c) 2008 The Thomson Corporation
 on

STN
 ACCESSION NUMBER: 1986:304474 BIOSIS <<LOGINID::20081202>>
 DOCUMENT NUMBER: PREV198682038380; BA82:38380
 TITLE: ONTOGENY OF THE ***PINEAPPLE*** ANANAS-COMOSUS
 SHOOT
 APEX.
 AUTHOR(S): MADHUSUDANAN K N [Reprint author]; NANDAKUMAR S
 CORPORATE SOURCE: DEP BOTANY, CALICUT UNIV, KERALA 673635
 SOURCE: Proceedings of the Indian National Science Academy
 Part B

376. Biological Sciences, (1985) Vol. 51, No. 3, pp. 369-

CODEN: PIBSBB. ISSN: 0073-6600.

DOCUMENT TYPE: Article

FILE SEGMENT: BA

LANGUAGE: ENGLISH

ENTRY DATE: Entered STN: 25 Jul 1986

Last Updated on STN: 25 Jul 1986

AB The morphological, histological and histochemical features of Ananas

comosus shoot apex were studied at seven stages of growth and development

under natural conditions. The stages selected were: the mature propagule

ready for planting (stage 1); two months after planting (stage 2); ten

months after planting (stage 3); 14 months after planting (stage 4); the

transitional or pre-floral stage (stage 5); the

organogenesis

stage (stage 6) and, the reversion stage of the inflorescence apex (stage

7). The apex of the propagule was characterized by a high protein content. The apex width, volume, cell population, and the protein content

decreased at stage 2; all these parameters were reversed at stage 3. The

nuclear area: cytoplasmic area, increased abruptly in the axial tunica

cells and central zone at stage 4, and decreased at stage 5. This ratio

increased in the lateral tunica and peripheral meristem in the evoked

stage (stage 5). The morphological, histological and histochemical features associated with the transition from stage 4 to stage 5,

under

natural conditions, resembled the changes noted under conditions of forced

flowering by the application of exogenous growth factors. The apex height, cell population and protein content decreased at stage 6.

The

apex at stage 7 resembled the one at stage 1 in many features but differed

in some.

=> FIL STNGUIDE

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

59.31

59.73

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE

TOTAL

ENTRY

SESSION

CA SUBSCRIBER PRICE

-3.20

-3.20

FILE 'STNGUIDE' ENTERED AT 10:59:08 ON 02 DEC 2008

USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT

COPYRIGHT (C) 2008 AMERICAN CHEMICAL SOCIETY (ACS)

FILE CONTAINS CURRENT INFORMATION.

LAST RELOADED: Nov 21, 2008 (20081121/UP).

=>

---Logging off of STN---

=>

Executing the logoff script...

=> LOG Y

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	1.50	61.23
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	0.00	-3.20

STN INTERNATIONAL LOGOFF AT 11:14:13 ON 02 DEC 2008